



To Reorder Call (281) 647-9700
or 1-800-295-BETA

Product Code: 5200-A



SUNSHINE

CITRUS SCENTED HOSPITAL SANITIZER, DISINFECTANT AND DEODORANT

FOR USE IN

- Hospitals • Ambulances • Bathrooms • Food Packaging Establishments •
- Locker Rooms • Athletic Mats • Urinals • Appliances • Toilets •
- Dental Clinics • Offices • Nursing Homes • Cafeterias • Industrial Plants • Stores •

DIRECTIONS

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This product is not to be used as a terminal sterilant/high level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the bloodstream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to preclean or decontaminate critical or semi-critical medical devices prior to sterilization or high level disinfection.

GENERAL: Shake container well before each use and hold it upright while spraying. For use on precleaned, hard nonporous surfaces; not for use on humans or animals. Not for use where food or related products are prepared, processed or stored. Remove food or related products from the area being treated. Avoid use where product would contact counters or tabletops where food may be stored, prepared or processed. Do not use on waxed/polished surfaces, water-based "latex" paint, shellac, cellulose (such as rayon) or other alcohol-sensitive materials.

To disinfect hard, non-porous surfaces: Preclean surfaces as directed above. Hold container upright about 6-8" from surface. Spray to thoroughly wet hard, nonporous surfaces. Treated surfaces must remain wet for 10 minutes. Allow to air dry. Rinse all surfaces that come in contact with food such as counters, appliances, tables with potable water. Do not use on utensils, glassware, and dishes.

Before use in federally inspected meat and poultry plants, all food products and packaging materials must be removed or carefully protected before disinfection.

To disinfect toilets and urinals: Remove gross filth prior to disinfection. Empty toilet bowl or urinal. Hold container upright about 6-8" from surface. Spray to thoroughly wet hard, nonporous surfaces. Brush over exposed surfaces and under the rim. Allow to stand for 10 minutes, then flush.

To disinfect Veterinary offices, animal areas, and animal laboratories: Remove all animals and feed from premises, vehicles, cages, kennels, etc. Remove all litter, droppings, and manure from floors, walls, and surfaces of facilities occupied or traversed by animals. Empty all feeding and water appliances. Thoroughly clean all surfaces with soap or detergent and rinse with water. Hold container upright about 6-8" from surface. Spray to thoroughly wet hard, nonporous surfaces. Treated surfaces must remain wet for 10 minutes. Allow to air dry. Immerse all halters, collars, leashes and other types of equipment used in handling and restraining animals as well as forks, shovels, and scrapers used for removing litter and manure. Ventilate buildings, vehicles, and other closed spaces. Do not house animals or employ equipment until treatment has been absorbed, set or dried. Thoroughly scrub feeding and watering equipment with soap or detergent, and rinse with potable water before reuse.

To Clean/Deodorize: Hold container upright about 6-8" from surface. Spray surface and allow to air dry.

To Control Mold/Mildew and Sanitize Porous Surfaces: Inhibits the growth of *Apergillus niger* and *Penicillium variable* fungi and controls *Staphylococcus aureus* and *Klebsiella pneumoniae* on porous surfaces such as fabric and leather surfaces. Hold container upright about 6-8" from surface. Spray to thoroughly wet surfaces. Treated surfaces must remain wet for 5 minutes. Allow surfaces to air dry. For mold/mildew control, repeat application every 7 days or more frequently if new growth appears.

TO KILL HIV-1 (Human Immunodeficiency virus Type 1; AIDS virus) ON PRECLEANED SURFACES/OBJECTS in health care or other settings with inanimate environmental surfaces/objects associated with the potential for transmission of HIV-1 and likely to be soiled with blood/body fluids, follow these additional **SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATING AGAINST HIV-1 ON SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS:** PERSONAL PROTECTION - Barrier protection items such as disposable latex gloves, gowns, masks, and eye coverings must be worn when handling items soiled with blood/body fluids. CLEANING PROCEDURE - Blood/body fluids must be thoroughly cleaned from surfaces/objects before application of this product as a disinfectant. CONTACT TIME - 10 minutes. DISPOSAL OF INFECTIOUS MATERIALS - Blood/body fluids and other materials removed from or used to clean surfaces/objects must be autoclaved and disposed of according to federal, state and local regulations for infectious waste disposal.

ACTIVE INGREDIENTS

2-Phenylphenol	0.20%
4-tert-Amylphenol	0.05%
Ethanol.....	53.000%

BETA DATA

FLASH POINT:	68°F Conc.	WETTING ABILITY:	None
PROPELLANT:	Hydrocarbon	FLAME EXTENSION:	12"
COLOR:	Clear	SOLUBILITY IN WATER:	100%
SPECIFIC GRAVITY:	0.88	PHOSPHATES:	None
TOXICITY:	1000 ppm OSHA TLV	BIODEGRADABILITY:	No
DETERGENCY:	None	pH:	7.2 Conc.
STORAGE STABILITY:	1 year +	% VOLATILE BY VOLUME:	99
COLD STABILITY:	0° F	EVAPORATION RATE:	Medium
ODOR:	Citrus	FORM:	Alcohol Base Aerosol

BETA TECHNOLOGY, INC.
CORPORATE HEADQUARTERS
HOUSTON, TX 77218